Research Journal of Pure Algebra -3(3), 2013, Page: 106-110 Available online through www.rjpa.info

PROPERTIES OF LATTICE ORDERED GROUPS AND ORDERED Γ - SEMIGROUPS

VB Subrahmanyeswara Rao Seetamraju*

Assistant Professor, VKR, VNB & AGK College of Engineering, Gudivada, Andhra Pradesh, India

(Received on: 20-02-13; Revised & Accepted on: 02-03-13)

ABSTRACT

In this paper, we present the properties of lattice ordered groups derived from the properties of partially ordered groups. The notion of Γ -semi groups was introduced by Sen in 1981. The concept of Γ - semigroups is a generalization of the concept of semigroups Many classical notions of semigroups have been extended to Γ -semigroups, (S, Γ, \leq) is called an ordered Γ -semigroup if S is a Γ -semigroup and (S, \leq) is a partially ordered set such that $a \leq b \Rightarrow a \gamma c \leq b \gamma c$ and $c \gamma a \leq c \gamma b$ for all $a, b, c \in S$ and $\gamma \in \Gamma$

AMS Subject classification (1991): 20M10

Keywords: Partially ordered groups, Lattice ordered groups, Semi groups, Γ - Semi groups, Ordered Γ - Semi groups.

1. INTRODUCTION

A semi group S is said to be partially ordered or a partially ordered semi group if it is associated with a partial ordering \leq which is defined by $a \leq b$ implies that $xay \leq xby$ for all x, y in S. The natural partial order which is an obvious partial ordering defined by $a \leq b$ if and only if a = cb for some $c = c^2 \epsilon S$. This natural partial ordering is compatible with multiplication. Some of the basic properties and results were given by Donald B. McAlister and some of the foundational results are due to A.H. Clifford. Suppose G is a partially ordered group i.e., G is a group partially ordered by \leq . Now $a \leq b$ if and only if $1 \leq a^{-1}b$ or equivalently $a \leq b$ if and only if $1 \leq ba^{-1}$

2. PROPERTIES OF PARTIALLY ORDERED, LATTICE ORDERED GROUPS

Here we consider the set G^+ consisting of elements exceeding the identity 1 and has the following properties:

(i) G^+ is a submonoid of G

(ii) a $G^+ = G^+ a$ for each a in G

(iii) 1 is the only invertible element of G^+

Notation 2.1: Let G be a partially ordered group and suppose that $a, b \in G$. Then the least upper bound of a and b is denoted by $a \lor b$, read as a join b. The greatest lower bound of a and b is denoted by $a \land b$, read as a meet b.

The following Proposition is due to Donald B.McAlister [2]

Proposition 2.2: Let G be a partially ordered group and suppose that $a, b \in G$. Then a and b have a least upper bound $a \lor b$ in G if and only if they have a greatest lower bound $a \land b$. This is only possible only when a^{-1} and b^{-1} have a least upper bound. In particular,

$a \wedge b = a(a \vee b)^{-1}b$ $a \wedge b = a(a \wedge b)^{-1}b$	$a \wedge b = (a^{-1} \vee b^{-1})^{-1}$
Also for any g in G ,	$u \lor b = (u \land b)$
$g(a \lor b) = ga \lor gb$ $(a \lor b)g = ag \lor bg$	$g(a \land b) = ga \land gb$ $(a \land b)g = ag \land b$

Corollary 2.3: *The following are equivalent for a partially ordered group G.*

(i) G is a \lor -semilattice under \leq

(ii) G is a \land -semilattice under \leq

(iii) $a \lor 1$ exists for each $a \in G$

(iv) $a \wedge 1$ exists for each $a \in G$

(v) $a \lor b$ exists for each $a, b \in G^+$

(vi) $a \wedge b$ exists for each $a, b \in G^+$

(vii) for each $a, b \in G^+$ there exists $c \in G^+$ such that $G^+a \wedge G^+b = G^+c$

Corresponding author: VB Subrahmanyeswara Rao Seetamraju Assistant Professor, VKR, VNB & AGK College of Engineering, Gudivada, Andhra Pradesh, India

VB Subrahmanyeswara Rao Seetamraju*/ PROPERTIES OF LATTICE ORDERED GROUPS AND ORDERED I- SEMIGROUPS/ RJPA- 3(3), March-2013.

If G satisfies one of the conditions in the above corollary, we say that G is a lattice ordered group or simply latticed group.

Definition 2.4: A lattice G is called a distributive lattice, if for any a, b, $c \in G$ (i) a \lor (b \land c) = (a \lor b) \land (a \lor c) and (ii) a \land (b \lor c) = (a \land b) \lor (a \land c). Clearly conditions (i) and (ii) are equivalent.

Definition 2.5: A lattice G is called a modular lattice, if for any $a, b, c \in G$ such that $a \le c$ implies $a \lor (b \land c) = (a \lor b) \land c$.

Theorem 2.6: Let G be a lattice ordered group under a partial order \leq , then G is a modular lattice under \leq .

Proof: The proof is obvious, because if G is a lattice ordered group with respect to a partial order \leq , then G is a distributive lattice under \leq and every distributive lattice is a modular lattice.

Definition 2.7: Two elements a, b of a lattice ordered group G are said to be orthogonal if $\mathbf{a} \wedge \mathbf{b} = \mathbf{1}$

Proposition 2.8: Let G be a lattice ordered group and let $a, b, c \in G$. If $a \wedge b = 1$, then $ac \wedge bc = c$.

Proof: Since $a \wedge b = 1$,

 $ac \wedge bc = 1 (ac \wedge bc)$ = $(a \wedge b) (ac \wedge bc)$ = $a (ac \wedge bc) \wedge b (ac \wedge bc)$ = $a^2c \wedge abc \wedge bac \wedge b^2 c$ = $a^2c \wedge abc \wedge b^2 c$, since $a \wedge a = a$ = $c (a(a \wedge b) \wedge b^2)$ = $c (a \wedge b^2)$ = $c, since a^m \wedge b^n = 1 \forall m, n \ge 0$

Definition 2.9: A homomorphism θ of a lattice ordered group G into a lattice ordered group H is an l-homomorphism if θ respects \vee and i.e., $(\mathbf{a} \vee \mathbf{b}) \theta = \mathbf{a}\theta \vee \mathbf{b}\theta$ and $(\mathbf{a} \wedge \mathbf{b}) \theta = \mathbf{a}\theta \wedge \mathbf{b}\theta$ for all $\mathbf{a}, \mathbf{b} \in \mathbf{G}$.

Definition 2.10: Let M and Γ be any two non-empty sets. If there exists a mapping $M \times \Gamma \times M \to M$, written (a, γ, b) by $a\gamma b$, M is called a Γ -Semigroup if M satisfies the identity $(a\gamma b)\mu c = a\gamma(b\mu c)$ for all $a, b, c \in M$ and $\gamma, \mu \in \Gamma$

Notation 2.11: Green's relations are defined on a Γ -Semigroup S as follows: $a\mathcal{L}b$ if and only if $S'\Gamma a = S'\Gamma b$ $a\mathcal{R}b$ if and only if $a\Gamma S' = b\Gamma S'$ $a\mathcal{D}b$ if and only if $a\Gamma\mathcal{L} \circ \mathcal{R}\Gamma b$ Where S'denotes S if S has an identity and $S' = S \cup \{1\}$ where 1 acts like the identity and $\mathcal{D} = \mathcal{R} \circ \mathcal{L}$

Definition 2.12: A semi group S is regular if and only if, for each $\mathbf{a} \in \mathbf{S}$, there exists $\mathbf{x} \in \mathbf{S}$ such that $\mathbf{a} = \mathbf{a}\alpha\mathbf{x}\alpha\mathbf{a}$ for $\alpha \in \Gamma$. In this case $\mathbf{a}' = \mathbf{x}\mathbf{a}\mathbf{x}$ is a solution to the pair of equations $\mathbf{a} = \mathbf{a}\mathbf{y}\mathbf{a}$, $\mathbf{y} = \mathbf{y}\mathbf{a}\mathbf{y}$. The element a is called an inverse for \mathbf{a} in S and the semi group is said to be an Inverse Γ - semi group if each element of S has a unique inverse in S.

3. GENERAL PROPERTIES OF Γ-SEMIGROUPS

Lemma 3.1: For every Γ – semigroup S, the following hold

- i) $a \leq e, a \in S, e \in E_s$ imply $a \in E_s$
- *ii*) $a \le b, b \in S, b$ regular imply a regular

iii) $a \le b, a, b \in S, a regular imply <math>a = eb = bf, e, f \in E_s, e \mathcal{D} f$

Proof: i) For $a \in S$, we have $a = x\alpha e = e\alpha y$ and $x\alpha a = a = a\alpha y$ for $x, y \in S', \alpha \in \Gamma$ Now, $a^2 = a\alpha a, \alpha \in \Gamma$ $= x\alpha e \alpha e\alpha y$ © 2013, RJPA. All Rights Reserved $= x\alpha e \alpha y$ $= a\alpha y$ = a $\therefore a \in E_s$

ii) Suppose $a \le b$, b regular then $a = x\alpha b = b\alpha y$; $\alpha = a\alpha y$

But for every inverse element b' of b, $a = a\alpha y = x\alpha b\alpha y$ = $x\alpha(b\alpha b'\alpha b)\alpha y$ = $(x\alpha b)\alpha b'\alpha b\alpha y$ = $a\alpha b'\alpha a$

Hence a is regular.

iii) Suppose $a \le b$, a is regular, which implies $a = x\alpha b = b\alpha y$ and $x\alpha a = a = a\alpha y(x, y \in S')$

But For every inverse element a' of a, $a = a\alpha a'\alpha a = a\alpha a'\alpha x\alpha b = e\alpha b$ for $e = a\alpha a'\alpha x$

Similarly, $a = b\alpha f$ for $f = y\alpha a'\beta a \in E_s$

Finally, since $e = a\alpha a'\beta x$, $a = e\alpha b$

 \therefore e \mathcal{R} a

Since $a = b\alpha f$, $f = y\alpha a'\beta a$, we have $a \mathcal{L} f$

Hence e \mathcal{D} f (:: $\mathcal{D} = \mathcal{R}$ o \mathcal{L})

Definition 3.2: (S, Γ, \leq) is called an ordered Γ – semigroup if (S, Γ) is a Γ –semigroup and (S, \leq) is a partially ordered set such that $a \leq b \Rightarrow a \gamma c \leq b \gamma c$ and $c \gamma a \leq c \gamma b$, for all $a, b, c \in S$ and $\gamma \in \Gamma$

The above property is similar to natural partial order compatibility with multiplication $a \le b \implies ac \le bc$ and $ca \le cb$ for all $c \in S$

Definition 3.3: The idempotents E_s of a Γ -semigroup can be given a partial order, ($E_s \neq \emptyset$) as follows: For e, $f \in E_s$ define $e \le f \iff e\alpha f = e = f\alpha e$

Lemma 3.4: The relation \leq is a partial order on E_s such that for $e, f \in E_s$ $e \leq f \iff e\alpha f = e = f\alpha e$

Proof: Firstly, for all $e \in E_s$, $e\alpha e = e$ Therefore $e \le e$ $\therefore \le$ is Reflexive

If $e \leq f$ and $f \leq e$, then $e = e\alpha f = f$

Therefore \leq is Anti-Symmetric

If $e \le f$ and $f \le h$, then $e = e\alpha f = e\alpha f\beta h = e\alpha h$ and $e = f\alpha e = h\alpha f\alpha e = h\alpha e$

Therefore $e \leq h$

Thus \leq is Transitive and hence the relation \leq is a partial order on E_s

Definition 3.5: If S is a commutative Γ – semigroup and all its elements are idempotents (S = E_s) then S is called a Γ –semilattice.

Hence for all $x, y \in S$: $x\alpha x = x$ and $x\alpha y = y\alpha x$

The relation \leq on idempotents is defined on the whole of a Γ – semilattice 'S'.

© 2013, RJPA. All Rights Reserved

VB Subrahmanyeswara Rao Seetamraju*/ PROPERTIES OF LATTICE ORDERED GROUPS AND ORDERED [- SEMIGROUPS/ RJPA- 3(3), March-2013.

An element g is a lower bound of elements e and f if $g \le e$ and $g \le f$ i.e., $g\alpha e = g = e\alpha g$ and $g\alpha f = g = f\alpha g$

Lemma 3.6: Let *S* be a Γ –semilattice. Then $e\alpha f \in S$ is the greatest lower bound of the elements *e* and *f* of *S*

Proof: Let $e\alpha f \in S$, $\alpha \in \Gamma$. Then $e\alpha f = f\alpha e = f\alpha f\alpha e = e\alpha f\alpha f$

 $\therefore (e\alpha f)\alpha f = e\alpha f = f\alpha(e\alpha f)$

This means $e\alpha f \leq f$

So, $(e\alpha f)\alpha e = f\alpha e\alpha e = f\alpha e = e\alpha f$

Therefore, $e\alpha(e\alpha f) = e\alpha(f\alpha e)$ = $e\alpha f\alpha e$ = $f\alpha e\alpha e$ = $f\alpha e$ = $e\alpha f$ $\therefore e\alpha f \leq e$

Thus $e\alpha f$ is a lower bound of e and f

If g is a lower bound of e and f, then

 $g = g\alpha f = g\alpha e$, and therefore $g\alpha(e\alpha f) = (g\alpha e)\alpha f = g\alpha f = g$ And hence $g \leq e\alpha f$

So, $e\alpha f \in S$ is the greatest lower bound of the elements *e* and *f*

4. PARTIAL ORDERING IN INVERSE Γ –SEMI GROUPS

In any Γ –semigroup S, the idempotent can be partially ordered by the relation $e \leq f \iff e\alpha f = e = f\alpha e \forall e, f \in S, \alpha \in \Gamma$

Analogous to this partial ordering, we can define a partial ordering to all the elements of an inverse Γ –semigroup by $x \le y \Leftrightarrow \exists e \in E_s : x = e\alpha y$

Since = $(x \alpha x^{-1})\alpha x$, $x\alpha x^{-1} \epsilon E_s$, we have \leq is reflexive

If $x = e\alpha y$ and $y = f\alpha x$, then

 $x = e\alpha y = e\alpha(e\alpha y) = e\alpha x$

Hence $x = e\alpha y = e\alpha f\alpha x = f\alpha x = y$, so \leq is antisymmetric

If $x = e\alpha y$ and $y = f\alpha z$, then

 $x = e\alpha y = e\alpha f\alpha z, e\alpha f \in E_s$, thus \leq is transitive

Lemma 4.1: In an inverse Γ – semigroup S, $x \leq y \iff x = x\alpha x^{-1}\alpha y$

Proof: $x \le y \Leftrightarrow \exists e \in E_s$: x = yae $\Leftrightarrow xax^{-1} = yax^{-1}$ $\Leftrightarrow x = xay^{-1}ax$ $\Leftrightarrow xax^{-1} = xay^{-1}$ $\Leftrightarrow x^{-1}ax = y^{-1}ax$ $\Leftrightarrow x = xax^{-1}ay$

Definition 4.2: Let S be an inverse Γ - semigrou. Then S is E-Unitary if and only if $e\alpha e = e = e\alpha a$ implies $a\alpha a = a$ for all $\alpha \in \Gamma$

Lemma 4.3: Let *S* be an *E*-Unitary inverse Γ – semigroup. Then $(x, y) \in \sigma$ if and only if $x\alpha x^{-1}\alpha y = y\alpha y^{-1}\alpha x$, σ is an equivalence on *S*

VB Subrahmanyeswara Rao Seetamraju*/ PROPERTIES OF LATTICE ORDERED GROUPS AND ORDERED [- SEMIGROUPS/ RJPA- 3(3), March-2013.

Proof: Suppose $x\alpha x^{-1}\alpha y = y\alpha y^{-1}\alpha x$. Then $e\alpha y = e\alpha x$ for $x, y \in S$ and $\alpha \in \Gamma$

$$\Rightarrow$$
 (*x*, *y*) $\in \sigma$

Conversely Suppose that $(x, y) \in \sigma$. Then $(x^{-1}, y^{-1}) \in \sigma$

 $\Rightarrow e\alpha x^{-1} = e\alpha y^{-1}$ for some $e \in E_s$

 $\Rightarrow e\alpha x^{-1}\alpha y = e\alpha y^{-1}\alpha y$ is idempotent

Since S is E-Unitary, $x^{-1}\alpha y$ is idempotent.

 $\therefore x^{-1} \alpha y = (x^{-1} \alpha y) \alpha (x^{-1} \alpha y)^{-1}$ $= x^{-1} \alpha y \alpha y^{-1} \alpha x$ $\Rightarrow x \alpha x^{-1} \alpha y = x \alpha x^{-1} \alpha y \alpha y^{-1} \alpha x = (y \alpha y^{-1}) \alpha (x \alpha x^{-1} \alpha x)$

= $y\alpha y^{-1}\alpha x$, Since idempotents commute.

ACKNOWLEDGMENTS

The author would like to thank the Management, Principal & Staff of VKR, VNB & AGK College Of Engineering, Gudivada for their valuable support and especially the Department of Mathematics.

REFERENCES

[1] A. H. Clifford And G.B. Preston, *Algebraic Theory of Semigroups*, Vol.1, 2, Math. Surveys No. 7, American Math. Soc., Providence, R.I., 1961, 1967.

[2] Donald B. Alister, *Lattice Ordered Groups and the Structure of Inverse Semigroups*, Notes for a course of lectures, University of Lisbon, May 1998.

[3] Heinz Mitsch, Semigroups and their natural order Math. Slovaca, 44(1994), No.4, 445-462.

[4] J.M. Howie, *Fundamentals of Semigroup Theory*, London Mathematical Society Monographs, New Series 12, Oxford, 1996.

[5] M.R. Darnel, Theory of Lattice Ordered Groups, Marcel Dekker Inc., New York, N.Y., 1995

[6] M.K, Sen and N.K. Saha, On Γ-semigroups I, Bull. Calcutta Math.Soc.78 (1986), No.3, 180-186

[7] M. Petrich, Introduction to semigroups, Merill, 1973.

Source of support: Nil, Conflict of interest: None Declared