

## THE EQUITABLE BONDAGE NUMBER OF A GRAPH

G. Deepak, N. D. Soner and Anwar Alwardi\*

Department of Studies in Mathematics, University of Mysore, Mysore-570 006, India

E-mail: [deepak1873@gmail.com](mailto:deepak1873@gmail.com), [ndsoner@yahoo.co.in](mailto:ndsoner@yahoo.co.in), [a\\_wardi@hotmail.com](mailto:a_wardi@hotmail.com)

(Received on: 25-12-11; Accepted on: 02-01-12)

### ABSTRACT

A subset  $D$  of  $V$  is called an equitable dominating set if for every  $v \in V - D$  there exists a vertex  $u \in D$  such that  $uv \in E(G)$  and  $|\deg(u) - \deg(v)| \leq 1$ . The minimum cardinality of such a dominating set is called the equitable domination number and is denoted by  $\gamma_e(G)$ . We define the equitable bondage number  $b_e(G)$  of a graph  $G$  to be the cardinality of a smallest set  $X \subseteq E$  of edges for which  $\gamma_e(G - X) > \gamma_e(G)$ . Sharp bounds are obtained for  $b_e(G)$  and the exact values are determined for some standard graphs.

**Keywords:** Graph, Bondage number, Equitable bondage number, Equitable domination number, Equitable domination.

**Mathematics Subject Classification :** 05C70.

### INTRODUCTION:

The graphs considered here are finite, undirected without loops and multiple edges having  $p$  vertices and  $q$  edges. Any undefined term in this paper may be found in Harary [3]. A set  $D$  of vertices in a graph  $G$  is a dominating set if each vertex of  $G$  that is not in  $D$  is adjacent to at least one vertex of  $D$ . The domination number  $\gamma(G)$  of  $G$  is the minimum cardinality of a dominating set of  $G$ . For a survey of results on domination (see [4]). A subset  $D$  of  $V$  is called an equitable dominating set if for every  $v \in V - D$  there exists a vertex  $u \in D$  such that  $uv \in E(G)$  and  $|\deg(u) - \deg(v)| \leq 1$ . The minimum cardinality of such a dominating set is denoted by  $\gamma_e(G)$  and is called the equitable domination number of  $G$ . The bondage number  $b(G)$  of  $G$  is the minimum cardinality among the sets of edges  $X \subseteq E$  such that  $\gamma(G - X) > \gamma(G)$  (see [2]). In this paper we now define the equitable bondage number of a graph  $G$ . The equitable bondage number  $b_e(G)$  of a graph  $G$  is the minimum cardinality of a set  $F \subseteq E$  of edges for which  $\gamma_e(G - F) > \gamma_e(G)$ .

### SOME EXACT VALUES:

In several instances we shall have cause to use the ceiling function of a number  $x$ ; that is denoted by  $\lceil x \rceil$  and takes the value of the least integer greater than or equal to  $x$ .

### RESULTS:

**Proposition 1:** The equitable bondage number of the complete graph  $K_p$  ( $p \geq 2$ ) is  $b_e(K_p) = \left\lceil \frac{p}{2} \right\rceil$ .

**Proof:** If  $H$  is a spanning subgraph of  $K_p$  that is obtained by removing fewer than  $\left\lceil \frac{p}{2} \right\rceil$  edges from  $K_p$ , then  $H$  contains a vertex of degree  $p - 1$ , whence  $\gamma_e(H) = 1$ . Thus  $b_e(K_p) \geq \left\lceil \frac{p}{2} \right\rceil$ .

We consider two cases.

**Case 1:** If  $p$  is even, the removal of  $\frac{p}{2}$  independent edges from  $K_p$  reduces the degree of each vertex to  $p - 2$  and therefore yields a graph  $H$  with equitable domination number  $\gamma_e(H) = 2$ .

\*Corresponding author: Anwar Alwardi\*, \*E-mail: [a\\_wardi@hotmail.com](mailto:a_wardi@hotmail.com)

**Case 2:** If  $p$  is odd, the removal of  $\frac{p-1}{2}$  independent edges from  $K_p$  leaves a graph having exactly one vertex of degree  $p-1$ , by removing one edge incident with this vertex, we obtain a graph  $H$  with  $\gamma_e(H) = 2$ . In both cases, the graph  $H$  resulted from the removal of  $\left\lceil \frac{p}{2} \right\rceil$  edges from  $K_p$ . Thus,  $b_e(K_p) = \left\lceil \frac{p}{2} \right\rceil$ .

We next determine the equitable bondage numbers of the cycle  $C_p$  and path  $P_p$  of order  $p$ .

**Lemma A[1]:** The equitable domination numbers of the cycle and path of order  $p$  are respectively

$$\begin{aligned}\gamma_e(C_p) &= \left\lceil \frac{p}{3} \right\rceil \text{ for } p \geq 3 \text{ and} \\ \gamma_e(P_p) &= \left\lceil \frac{p}{3} \right\rceil \text{ for } p \geq 1.\end{aligned}$$

**Theorem 2:** Let  $K_{m,n}$  be a complete bipartite graph with  $|m-n| \leq 1$  and  $m \leq n$  then  $b_e(K_{m,n}) = m$ .

**Proof :** Let  $V = V_1 \cup V_2$  be the vertex set of  $K_{m,n}$  where  $|V_1| = m$  and  $|V_2| = n$ . Let  $v \in V_2$  then by removing all edges incident with  $v$  we obtain a graph  $H$  containing two components  $K_1$  and  $K_{m,n-1}$ . Hence

$$\begin{aligned}\gamma_e(H) &= \gamma_e(K_1) + \gamma_e(K_{m,n-1}) \\ &= 1 + \gamma_e(K_{m,n}) \\ &> \gamma_e(K_{m,n})\end{aligned}$$

Thus,

$$\begin{aligned}b_e(K_{m,n}) &= \deg(v) \\ &= |V_1| \\ &= m.\end{aligned}$$

**Theorem 3:** The equitable bondage number of the  $p$ -cycle is

$$b_e(C_p) = \begin{cases} 3 & \text{if } p \equiv 1 \pmod{3} \\ 2 & \text{otherwise.} \end{cases}$$

**Proof:** Since  $\gamma_e(C_p) = \gamma_e(P_p)$  for  $p \geq 3$ , we see that  $b_e(C_p) \geq 2$ . If  $p \equiv 1 \pmod{3}$  the removal of two edges from  $C_p$  leaves a graph  $H$  consisting of two paths  $X$  and  $Y$ . If  $X$  has order  $p_1$  and  $Y$  has order  $p_2$  then either  $p_1 \equiv p_2 \equiv 2 \pmod{3}$  or without loss of generality,  $p_1 \equiv 0 \pmod{3}$  and  $p_2 \equiv 1 \pmod{3}$ . In the former case,

$$\begin{aligned}\gamma_e(H) &= \gamma_e(X) + \gamma_e(Y) = \left\lceil \frac{p_1}{3} \right\rceil + \left\lceil \frac{p_2}{3} \right\rceil \\ &= \frac{(p_1+1)}{3} + \frac{(p_2+1)}{3} \\ &= \frac{p+2}{3} > \left\lceil \frac{p}{3} \right\rceil = \gamma_e(C_p)\end{aligned}$$

In the latter case,

$$\begin{aligned}\gamma_e(H) &= \frac{p_1}{3} + \frac{(p_2+2)}{3} \\ &= \frac{(p+2)}{3} > \left\lceil \frac{p}{3} \right\rceil = \gamma_e(C_p).\end{aligned}$$

In either case, when  $p \equiv 1 \pmod{3}$ , we have  $\gamma_e(C_p) \geq 3$ . To obtain the upper bounds that by trichotomy, will yield the desired equalities of our theorem's statement we consider two cases.

**Case 1:** Suppose that  $p \equiv 0, 2 \pmod{3}$ . The graph  $H$  obtained by removing two adjacent edges from  $C_p$  consists of an equitable isolated vertex and a path of order  $p-1$ . Thus,

$$\gamma_e(H) = 1 + \gamma_e(P_{p-1}) = 1 + \left\lceil \frac{p-1}{3} \right\rceil$$

$$= 1 + \left\lceil \frac{p}{3} \right\rceil > \left\lceil \frac{p}{3} \right\rceil = \gamma_e(C_p)$$

hence  $b_e(C_p) \leq 2$  in this case. Combining this with the upper bound obtained earlier, we have  $b_e(C_p) = 2$  if  $p \equiv 0, 2 \pmod{3}$ .

**Case 2:** Suppose now that  $p \equiv 1 \pmod{3}$ . The graph H resulting from the deletion of three consecutive edges of  $C_p$  consists of two equitable isolated vertices and a path of order  $n - 2$ . Thus,

$$\begin{aligned} \gamma_e(H) &= 2 + \left\lceil \frac{(p-2)}{3} \right\rceil = 2 + \frac{(p-1)}{3} \\ &= 2 + \left\lceil \frac{p}{3} \right\rceil - 1 > \left\lceil \frac{p}{3} \right\rceil = \gamma_e(C_p) \end{aligned}$$

so that  $b_e(C_p) \leq 3$ . With the earlier inequality, we conclude that  $b_e(C_p) = 3$  when  $p \equiv 1 \pmod{3}$ .

As an immediate corollary to the Theorem 3 we have the following

**Corollary 3.1:** The equitable bondage number of the path of order  $(p \geq 2)$  is given by

$$b_e(P_p) = \begin{cases} 2 & \text{if } p \equiv 1 \pmod{3} \\ 1 & \text{otherwise.} \end{cases}$$

**Theorem 4:** The equitable bondage number of the wheel  $W_p$  is

$$b_e(W_p) = \begin{cases} 3 & \text{if } p \equiv 2 \pmod{3} \\ 2 & \text{otherwise.} \end{cases}$$

**Proof:** Let  $W_p = K_1 + C_{p-1}$  and label  $C_{p-1}$ :  $e_1, e_2, \dots, e_{p-1}$  be the edges of  $C_{p-1}$ . We consider two cases.

**Case 1:** Suppose that  $p \equiv 2 \pmod{3}$ . The graph H obtained by removing three consecutive edges  $e_1, e_2, e_3$  on  $C_{p-1}$  form  $W_p$  consists of three equitable isolated vertices and a path of order  $p - 3$ . Thus,

$$\begin{aligned} \gamma_e(H) &= 3 + \gamma_e(P_{p-3}) \\ &= 3 + \left\lceil \frac{p-3}{3} \right\rceil > \gamma_e(W_p) \end{aligned}$$

Hence,  $b_e(W_p) = 3$ .

**Case 2:** Suppose that  $p \not\equiv 2 \pmod{3}$ . The graph H resulting from the deletion of two adjacent edges  $e_1$  and  $e_2$  on  $C_{p-1}$  from  $W_p$  consists of two equitable isolated vertices and a path of order  $p - 2$ . Thus,

$$\begin{aligned} \gamma_e(H) &= 2 + \gamma_e(P_{p-2}) \\ &= 2 + \left\lceil \frac{p-2}{3} \right\rceil > \gamma_e(W_p) \end{aligned}$$

Hence,  $b_e(W_p) = 2$ .

**Theorem 5:** If T is non trivial tree, then  $b_e(T) \leq 2$ .

**Proof:** If T is of order 2 or 3 then it is clear that  $b_e(T) = 1$ . We are interested here about the equitable edge in the tree. Suppose that T has at least four vertices, then we can classify trees into 2 cases.

**Case 1:** According to the end edge of the trees, if all the end edge are not equitable then by deleting any equitable edge the equitable domination number will be increasing i.e.,  $b_e(T) = 1$ .

**Case 2:** If the end edge are equitable then we have two sub cases, if there exists leaf which is  $P_4$  then  $b_e(T) = 2$  otherwise  $b_e(T) = 1$ .

Hence for any nontrivial tree then  $b_e(T) \leq 2$ .

**Theorem 6:** If  $G$  be any graph and  $x$  be equitable edge then  $\gamma(G - x) \geq \gamma(G)$ .

**Proof:** Since we have a theorem, for any edge  $x$  of a graph,  $\gamma(G - x) \geq \gamma(G)$  and from the definition of equitable domination number if the graph without any equitable isolated vertex then the  $\gamma_e(G) = \gamma(G)$  and of course if we delete any equitable edge  $x$  from the graph  $G$  we get  $\gamma(G - x) \geq \gamma(G)$ .

**Theorem 7:** If  $G$  is a connected graph of order  $p \geq 2$ , then  $b_e(G) \leq p - 1$ .

**Proof :** Let  $u$  and  $v$  be adjacent vertices with  $|\deg(u) - \deg(v)| \leq 1$  and  $\deg(u) \leq \deg(v)$ . Let  $E_u$  denote the set of edges incident with  $u$ . Then  $\gamma_e(G - E_u) = \gamma_e(u)$  and  $\gamma_e(G - u) = \gamma_e(G) - 1$ . Also, if  $D$  denotes the union of all minimum equitable dominating sets for  $G - u$ , then  $u$  is adjacent in  $G$  to no vertex of  $D$ . Hence  $|E_u| \leq p - 1 - |D|$  and  $u \notin D$ . Now if  $F_v$  denotes the set of edges from  $v$  to a vertex in  $D$ , then since  $v \notin D$  we must have

$$\begin{aligned} \gamma_e(G - u - F_v) &> \gamma_e(G - u) \text{ or equivalently,} \\ \gamma_e(G - u - F_v) &> \gamma_e(G) - 1 \end{aligned}$$

Thus,  $\gamma_e(G - (E_u \cup F_v)) > \gamma_e(G)$  and

$$\begin{aligned} \text{we see that, } b_e(G) &\leq |E_u \cup F_v| \\ &= |E_u| + |F_v| \\ &\leq (p - 1 - |D|) + |D| \\ &= p - 1 \end{aligned}$$

#### REFERENCES:

- [1] K. D. Dharmalingam, Studies in Graph Theory – Equitable Domination and bottleneck domination, Ph.D. Thesis (2006).
- [2] J. F. Fink, M.S. Jacobson, L. F. Kinch and J. Roberts. The bondage number of a graph. Discrete Math. 86 (1990) 47-57.
- [3] F. Harary, Graph Theory, Addison-Wessley, Reading Mass. (1969).
- [4] T. W. Haynes, S. T. Hedetniemi and P.J. Slater, Fundamentals of domination in graphs, Marcel Dekker, Inc., New York (1998).

\*\*\*\*\*